

Ariunsuren, P., D. Damiran, and P. Stevens. 2009. Results of assessment of the current status of degree of disturbance of the desert-steppe rangelands of western Mongolia. In: Proc. Soils & Crops conference. 25-26 February, 2009. Saskatoon, SK; Canada.

Results of Assessment of the Current Status and Degree of Disturbance of the Desert-Steppe Rangelands of Western Mongolia

P. Ariunsuren¹, D. Damiran² and J.P. Stevens²

¹Department of Biology, Mongolian State University of Agriculture, Ulaanbaatar, Mongolia, 210153

²Department of Animal Science and Poultry Science, University of Saskatchewan, Saskatoon, SK, S7N 5A8

Key Words: desertification, invasion, Mongolian rangelands, soil erosion, sustainability

Abstract

Concern about the status of Mongolia's rangelands has increased due to dramatic increases in livestock populations of this country during the past 18 years. The objective of this study was to document the current status and degree of disturbance of the desert-steppe rangelands of western Mongolia. A total of 123 rangeland monitoring sites representing 33 million hectares (21 % of total country area) were established and inventoried in July/August 2008. Overall, our study indicated that 82.9% of the desert-steppe rangelands, consisting of 27.36 million hectares, have been degraded to some extent. The degree of degradation varies; slight 26%, moderate 30.9%, severe 13.8% and very severe 12.2% of total the area. In conclusion, in addition to natural factors several specific human-driven causes, such as animal husbandry have led to widespread disturbance in Mongolian desert-steppe rangelands.

Introduction

Mongolia is a land-locked country located between 87°41" and 119°56" east longitude, and 41°35" and 52°09" north latitude in the central region of north-east Asia (Tsegmid and Vorobiev, 1999). It has a population of 2.6 million people in an area of 1.56 million km², with considerable potential for agricultural production, primarily through extensive livestock pastoral grazing. The territory of Mongolia as a whole is characterized by great diversity and can be separated into several agro-ecological regions and sub-regions with significantly different environments and vegetation cover which provide different agricultural and livestock production potentials (Jigjidsuren and Johnson, 2003). The main ecological regions are: Hangai-Hovsgul, Selenge-Onon, Central and Eastern Steppe, Gobi Desert Region, and Mongolian Altai Region (Johnson et al., 2006). The Altai Region, located in western Mongolia includes two districts. The first includes the Mongolian Altai stretching from the north-west to the south-east for more than 1500 km. The second district comprises the Turgen Mountains and Lake Ureg-Nur. The Altai Region's elevation ranges between 1500 and 4000 m (Tsegmid and Vorobiev, 1990). The mean annual temperature is between -2.5°C and 5.0°C with low temperature of -24°C in January and high temperature of 22°C in July; between 60-120 frost free days, and precipitation between 400 and 500 mm (Batjargal, 1996). Snow depth ranges between 5 to >15 mm and wind speed can occur between 2 and 6 m/sec (Tsegmid and Vorobiev, 1990). The Mongolian Altai north-eastern

Ariunsuren, P., D. Damiran, and P. Stevens. 2009. Results of assessment of the current status of degree of disturbance of the desert-steppe rangelands of western Mongolia. In: Proc. Soils & Crops conference. 25-26 February, 2009. Saskatoon, SK; Canada.

slopes have a desert steppe and desert ecosystem (western Mongolian desert steppe) (Gunin, 1999).

Western Mongolian desert-steppe rangelands play an important role in the Mongolian economy. An annual precipitation ranging from less 50 mm in the south to over 150 mm at upper elevations in the west creates an arid to semiarid shrub-dominated environment. This gradient generates correspondingly large differences in plant production. Concern about the status of Mongolia's rangelands has increased due to dramatic increases in livestock populations during the past 18 years. Between 1990 and 2007, the number of domestic animals increased to an estimated 43.3 million head of livestock (Monthly Bulletin of Statistics, 2008.). An overall decline in net primary productivity has been observed with increasing signs of localized degradation, including both plant community changes and increased soil erosion. The aim of this study was to document the current status and degree of disturbance of the western Mongolian desert-steppe rangelands. Our specific research objectives were 1) to document plant communities sampled along gradients of desert-steppe zones of western Mongolia, 2) evaluate current rangeland conditions, 3) to establish permanent monitoring sites. A further objective was to develop reference site descriptions in the desert-steppe rangelands of the western Mongolia.

Materials and Methods

The study area was the desert steppe ecosystem type of western Mongolia, in Khovd and Uvs Aimag (province). This ecosystem is the largest rangeland type in this area (Buyan-Orshih, 1981; Gunin, 1999). Based on the environmental factors of slope, elevation, soil type, or soil moisture content, and vegetation, we stratified the sampling area into 5 habitats: 1) depression, 2) lower slopes of a mountain (lower slopes), 3) river flood plains, 4) mountain steppe and 5) valley. Permanent monitoring sites were established and a survey was conducted from mid July to late August 2008. In total 123 monitoring sites were spread throughout the 5 habitats (Fig. 1). We established and sampled 18, 10, 56, 18, and 21 monitoring sites for depression, lower slopes, river flood plains, mountain steppe, and valley habitat, respectively.

Establishing Monitoring Sites

In the selection of monitoring sites, local (soum and bag) governors and herders were consulted to determine the most appropriate resource management area for the survey. We also considered herders' mobility and camp locations in the research area. We were able to cover the pastures for all seasons, including winter, transitional or spring/autumn, and summer ranges. These permanent monitoring sites were spread between 91°18'23.1" and 93°21'49.0" of eastern longitude, and 45°06'49.0" and 50°46'41.8" of the northern latitude. All monitoring sites were marked with the geographical positions enabling investigators to use them as long term permanent monitoring sites, as well as for short and mid term research purposes. Each monitoring site was permanently marked using stakes and documented with photographs.

Rangeland Survey

At each monitoring site, three sampling plots were oriented along cardinal directions from the center of the monitoring point (25 meter paces each direction). The vegetation was surveyed using the Braun-Blanquet cover scale: (+) a limited number of individuals; (1) vegetation cover < 5 %; (2) vegetation cover between 5 and 25 %; (3) vegetation cover between 25 and 50 %; (4)

Ariunsuren, P., D. Damiran, and P. Stevens. 2009. Results of assessment of the current status of degree of disturbance of the desert-steppe rangelands of western Mongolia. In: Proc. Soils & Crops conference. 25-26 February, 2009. Saskatoon, SK; Canada.

In the area, where this study was conducted, established reference sites or detailed description of reference sites were lacking. Therefore published materials (Gubanov, 1996; Gunin et al., 1999; Jigjidsuren and Johnson, 2003; Tserendash and Altanzul, 2006) of related studies were the main references to develop reference sites or indicator species from healthy ecotypes, as well as our observations from previous and current studies (Damiran, 2005; Ariunsuren, 2006). Indicator plant species for healthy ranges were selected as follows:

Depression: Selected species or key species for six functional groups included *Eurotia ceratoides* (L.) C. A. Mey., *Suaeda kossinskyi* Iljin. or *Suaeda heterophylla* (Kar. et Kir.) Bge., *Reaumuria soongorica* (Pall.) Maxim. or *Salsola* spp., *Agrophyron cristatum* (L.) P. B., *Nanophyton grubovii* Prato., *Cleistogenes squarrosa* (Trin.) Keng. and *Caragana* spp. Healthy sites vegetation canopy cover ranges from 41-50%.

Lower slopes: Healthy site key species are: *Stipa Krylovii* Roshev., *Agrophyron cristatum* (L.) P. B. or *Koeleria cristata* (L.) Pers., *Artemisia frigida* Willd., *Potentilla* spp., *Ephedra* spp. *Heteropappus altaicus* (Wiild.) Novopokr. and *Convolvulus Ammanii* Desr. dominated. Canopy coverage is 41-50%.

River flood plains: In the river flood plains of desertified steppes, the fine succession series are dominated by *Carex duruiscula* C. A. Mey., *Halerpestes salsuginosa* (Pall ex Georgi.) Greene., *Glaux maritima* L., *Saussurea amara* (L.) DC., *Plantago salsa* Pall., *Oxytropis salina* Vass., *Iris lacteal* Pall., and *Suaeda corniculata* (C. A. Mey.) Bge.. Rangeland sites vegetation canopy cover could range between 81 to 100%.

Mountain steppe: The mountain steppe is dominated by crested wheatgrass (*Agrophyron cristatum* (L.) P. B.), *Stipa Krylovii* Roshev., *Festuca lenensis* Drob., *Poa attenuata* Trin., Junegrass (*Koeleria cristata* (L.) Pers.) and *Carex pediformis* C.A. Mey. (Bedunah, Schmidt, 2000). Further, forb-sedge (*Carex duruiscula* C. A. Mey. and *C. stenophylloides* V. Krecz.), grass-forb (*Agrophyron cristatum* (L.) P. B., *Stipa Krylovii* Roshev., *Poa attenuata* Trin., *Poa altaica* Trin., *Koeleria cristata* (L.) Pers., *Koeleria altaica* (Domin.) Kryl.), and fescue mountain steppe (*Festuca lenensis* Drob.) are the dominant communities in this habitat. In the mountain steppe, the expected level of canopy cover would be in the range of 81 to 100%.

Valley: In the valley of western Mongolian desert steppe *Anabasis brevifolia* C.A.Mey, *Allium polyrrhizum* Turcz. ex Rgl., *Stipa gobica* Roshev., *Salsola passerina* Bge., *Reaumuria soongorica* (Pall.) Maxim., and *Ajania fruticulosa* (Ledeb.) Poljak. are key plant species. Vegetation cover of healthy rangelands should be in the range 21-25%.

Results and Discussion

During the 2008 survey, a total of 110 species which belong to 60 genera, and 22 families were identified among the monitoring sites. In total, 41 plant community types were distinguished. The standing biomass range was 2.4-33.8 kg ha⁻¹, 3.6-34.1 kg ha⁻¹, 57-93 kg ha⁻¹, 2.4-14.3 kg ha⁻¹, and 2-38.0 kg ha⁻¹ for the depression, lower slopes, river flood plains, mountain steppe, and valley habitat respectively. An average number of plant species were 6, 5, 4, 5, and 4 for the depression, lower slopes, river flood plains, mountain steppe, and valley habitat respectively. Based on the current study, a range of vegetation canopy coverage by rangeland health category and by habitat in desert steppe of western Mongolia is reported (Table 1). This study also found that 16 monitoring sites were in a healthy condition. Whereas 40, 35, 21, and 11 monitoring sites are slight, moderate, severe, and very severely disturbed, respectively. Habitats differed in vegetation canopy cover between sites for different rangeland health categories.

Ariunsuren, P., D. Damiran, and P. Stevens. 2009. Results of assessment of the current status of degree of disturbance of the desert-steppe rangelands of western Mongolia. In: Proc. Soils & Crops conference. 25-26 February, 2009. Saskatoon, SK; Canada.

Table 1. Vegetation Canopy Coverage (%) by 6 Habitats of the Desert-steppe Rangelands of Western Mongolia

Rangeland Health Categories	Habitats					
	Depression	Flood Plains		Valley	Lower Slope	Mountain Steppe
		river plain	salt-marsh			
Healthy	41-50	81-100	41-50	21-25	41-50	81-100
Disturbed						
Slight	31-40	60-80	31-40	16-20	31-40	61-80
Moderate	21-30	40-59	21-30	11-15	21-30	41-60
Severe	11-20	20-39	11-20	6-10	11-20	21-40
Very severe	0-10	0-19	0-10	0-5	0-10	0-20

Variation in the rangeland disturbance level was found between habitats (Table 2). The most pronounced degrading of vegetation succession in the flood plains of western Mongolia are caused by excessive grazing. Hydromorphic vegetation succession changes can be observed in the desert steppe when small ponds are drying up and turning into meadow saline soil depressions. In the most hydrophilous reed stands (*Achnatherum splendens* + *Phragmites australis*), the shore developed a halohydromorphic community with *Achnatherum splendens* (Trin.) Nevski., *Phragmites australis* (Cav.) Trin ex Steud., *Hordeum Bogdanii* Wilensky., *Puccinella Hauptiana* V. (Krecz.) Kitag., *Halerpestes salsuginosa* (Pall. ex Georgi.) Greene., *Oxytropis salina* Vass. and *Potentilla* spp. This survey revealed that the most pronounced degrading of vegetation succession in the flood plains of western Mongolia is caused by excessive grazing.

Table 2. Rangeland Disturbance Level in Six Habitats of the Desert-steppe Rangelands of Western Mongolia

Rangeland Health Categories ¹	Habitats					
	Depression	Flood Plains		Valley	Lower Slope	Mountain Steppe
		river plain	salt-marsh			
Healthy	1	2	2	5	2	4
Disturbed						
Slight	8	1	1	13	7	10
Moderate	6	2	0	18	3	6
Severe	1	1	1	12	5	1
Very severe	2	-	-	8	1	-
<i>n</i>	18	6	4	56	18	21

¹Rangeland health categories based on similarity index (Oosting, 1956) of dominant species to reference site indicator species and vegetation inventories surveyed by Braun-Blanquet, (1932) methodology. If similarity index was 100-80, 80-67, 67-50, 50-29, 29.0-0.0, current status of sites were considered as normal, slight, moderate, severe, and very severely disturbed.

Conclusions and Implications

In conclusion, a combination of natural factors and several specific human-driven factors, including vehicular traffic and especially animal husbandry, are the driving forces for widespread disturbance in dessert-steppe rangelands of western Mongolia. Established permanent monitoring sites will provide reference material and a benchmark of the desert-steppe

Ariunsuren, P., D. Damiran, and P. Stevens. 2009. Results of assessment of the current status of degree of disturbance of the desert-steppe rangelands of western Mongolia. In: Proc. Soils & Crops conference. 25-26 February, 2009. Saskatoon, SK; Canada.

rangeland of western Mongolia for future investigators and managers. Revisiting past studies for comparative analysis will provide information useful for establishing pastureland ecological conditions and trends. This is especially important in the desert steppe of western Mongolia where significant changes are happening and may soon accelerate due to the development of mineral resources.

Acknowledgements

The project was supported by the Training for Rural Development Project, a Canadian International Development Agency (CIDA) project administered by the University of Saskatchewan, Canada and Mongolian State University Agriculture, Mongolia. We thank B. Battulga, E. Erkhembayr, D. Ser-Od, and E. Ulziibayr for their assistance with the field work.

References

- Adams, B.W., G., Ehlert, C. Stone, M. Alexander, D. Lawrence, M. Willoughby, D. Moisey, C. Hincz, A. Burkinshaw. 2005. Range health assessment for grassland, forest and tame pasture. Public Lands and Forest Division, Alberta Sustainable Resources Development. Pub. No. T/044.
- Ariunsuren, P. 2006. Study on *Poa* L. species in Mongolia and Inner Mongolia in terms of taxonomy, geography and ecology. Proceedings of an international workshop on Participation by Researchers for Ecological problems. Ulaanbaatar. 29-33 p.
- Batjargal, Zh (ed.). 1996. Nature and environment in Mongolia (in Mongolian). Ministry of nature and the environment. Ulaanbaatar. 131 pp.
- Braun-Blanquet, J. 1932. Plant sociology: The Study of plant communities. Translated by G.D. Fuller and H.S. Conard. McGraw Hill Book Co., New York, NY. 430 pp.
- Buyan-Orshikh, Kh. 1981. Flora and vegetation of Sandy plant of the Great lakes depression. Ulaanbaatar: Press of Mongolian Academy of Science (*in Mongolian*). 297 pp.
- Damiran, D. 2005. Palatability of Mongolian Rangeland Plants. Circular of Information No.3. Union, Oregon, USA: Oregon State University, Eastern Oregon Agricultural Research Station. 92 p.
- Gunin P.D., E.A. Vostokova, N.I. Dorofeyuk, P.E. Tarasov, C.C. Black. 1999. Vegetation dynamics of Mongolia. Dordrecht: Kluwer. 238 pp
- Jigjidsuren, S., and D.A. Johnson. 2003. Forage plants in Mongolia. Admon Press. Ulaanbaatar. 503 pp.
- Johnson, D. A., D. P. Sheehy, D. Miller, D. Damiran. 2006. Mongolian rangelands in transition. J. Sècheresse 17(1):133-141.
- Monthly Bulletin of Statistics. 2008. Mongolian statistics book. Ulaanbaatar, Mongolia: Printed by the National Statistical Office. 144 pp.
- Oosting, H.J., 1956. The study of plant communities. San Francisco, CA: W.H. Freeman and Company. 440 pp.
- Tsegmid, Sh., V.V. Vorobiev (eds.). 1990. The national atlas. Mongolian People's Republic (in Mongolian). GUGK SSSR, GUGK MNR, Moscow, Ulaanbaatar, 144 pp.
- Tserendash, S., Ts. Altanzul. 2006. Guide booklet of grazing management. [*in Mongolian*]. Ulaanbaatar, Mongolia: EDO Publishing Co., ltd. 52 pp.
- Ulziikhutag, N (ed.). 1984. Dictionary of terminology of Mongolian plants (in Mongolian). State Publishing House, Ulaanbaatar. 360 pp.

Ariunsuren, P., D. Damiran, and P. Stevens. 2009. Results of assessment of the current status of degree of disturbance of the desert-steppe rangelands of western Mongolia. In: Proc. Soils & Crops conference. 25-26 February, 2009. Saskatoon, SK; Canada.